



Cultivation of better starting material with cuttings dipped in ArgicinPlus

More homogeneous cuttings with ArgicinPlus

In recent years we have gained experience at various strawberry propagators. After immersing the strawberry cutting into ArgicinPlus, the cutting takes root faster and the final tray plants are more homogeneous. This applies to both June-bearing and in everbearing strawberries.

In a number of batches this resulted in a difference of up to 40% in rooting and continued growth.

Use ArgicinPlus when immersing cuttings:

- 1 Litre ArgicinPlus in 500 Litres of water
- immerse the cutting for 15 minutes
- leave to leak and then plant or pot

This gives your cutting or tray plant time to sufficiently absorb substances from the willow bark extract in the ArgicinPlus. These substances make your crop more resilient:

- by increasing the robustness and regenerative capacity of your crop, and
- with extra callus formation, which in turn leads to more roots



Left hand side

Row of trays where the cuttings have been immersed in a solution containing ArgicinPlus.

Right hand side

Untreated

Conclusion

With ArgicinPlus: less waste in the production of tray plants and waiting-bed plants, more robust and therefore better plants for the production grower.

**TIP**

We also have advice for **production growers** based both on our experience and elaborate testing.

Get to know our products

PlantoSys develops products to make and keep crops resilient, robust and vital. And with results.

For more information, please visit www.plantosys.com/en